

# Package ‘ArDec’

February 19, 2015

**Type** Package

**Title** Time series autoregressive-based decomposition

**Version** 2.0

**Date** 2013-05-09

**Author** Susana Barbosa

**Maintainer** Susana Barbosa <[sabarbosa@fc.ul.pt](mailto:sabarbosa@fc.ul.pt)>

**Description** Package ArDec implements autoregressive-based decomposition of a time series based on the constructive approach in West (1997). Particular cases include the extraction of trend and seasonal components.

**License** GPL (>= 2)

**NeedsCompilation** no

**Repository** CRAN

**Date/Publication** 2013-05-10 10:03:26

## R topics documented:

ardec . . . . .	1
ardec.lm . . . . .	3
ardec.periodic . . . . .	4
ardec.trend . . . . .	5
tempEng . . . . .	6

## Index

7

---

ardec *Time series autoregressive decomposition*

---

## Description

Decomposition of a time series into latent subseries from a fitted autoregressive model

**Usage**

```
ardec(x, coef, ...)
```

**Arguments**

x	time series
coef	autoregressive parameters of AR(p) model
...	additional arguments for specific methods

**Details**

If an observed time series can be adequately described by an (eventually high order) autoregressive AR( $p$ ) process, a constructive result (West, 1997) yields a time series decomposition in terms of latent components following either AR(1) or AR(2) processes depending on the eigenvalues of the state evolution matrix.

Complex eigenvalues  $r \exp(iw)$  correspond to pseudo-periodic oscillations as a damped sine wave with fixed period ( $2\pi/w$ ) and damping factor  $r$ . Real eigenvalues correspond to a first order autoregressive process with parameter  $r$ .

**Value**

A list with components:

period	periods of latent components
modulus	damping factors of latent components
comps	matrix of latent components

**Author(s)**

S. M. Barbosa

**References**

West, M. (1997), Time series decomposition. Biometrika, 84, 489-494.

West, M. and Harrison, P.J. (1997), Bayesian Forecasting and Dynamic Models, Springer-Verlag.

**Examples**

```
data(tempEng)
coef=ardec.lm(tempEng)$coefficients

# warning: running the next command can be time consuming!

decomposition=ardec(tempEng,coef)
```

---

**ardec.lm***Fit an autoregressive model as a linear regression*

---

## Description

Function ardec.lm fits an autoregressive model of order p, AR(p) to a time series through a linear least squares regression.

## Usage

```
ardec.lm(x)
```

## Arguments

x                   time series

## Value

For ardec.lm, an object of class "lm".

## Author(s)

S. M. Barbosa

## References

West, M. (1995), Bayesian inference in cyclical component dynamic linear models.Journal of the American Statistical Association, 90, 1301-1312.

## See Also

[ar](#), [lm](#)

## Examples

```
data(tempEng)
model=ardec.lm(tempEng)
```

---

ardec.periodic	<i>Extraction of individual periodic components from a monthly time series</i>
----------------	--

---

## Description

Function ardec.periodic extracts a periodic component from the autoregressive decomposition of a monthly time series.

## Usage

```
ardec.periodic(x, per, tol = 0.95)
```

## Arguments

x	time series
per	period of the component to be extracted
tol	tolerance for the period of the component

## Value

A list with components:

period	period for the annual component
modulus	damping factor for the annual component
component	extracted component

## Author(s)

S. M. Barbosa

## Examples

```
data(tempEng)
ardec.periodic(tempEng,per=12)
```

---

**ardec.trend***Estimation of the trend component from a monthly time series*

---

## Description

Function ardec.trend extracts the trend component from the autoregressive decomposition of a monthly time series.

## Usage

```
ardec.trend(x)
```

## Arguments

x time series

## Value

A list with components:

modulus	damping factor for the annual component
trend	trend component

## Author(s)

S. M. Barbosa

## Examples

```
data(co2)
ardec.trend(co2)
```

---

tempEng	<i>Time series of monthly temperature values</i>
---------	--

---

**Description**

Monthly temperature in Central England from 1723-1970

**Usage**

```
data(tempEng)
```

**Format**

Time-Series [1:2976] from 1723 to 1971

**Source**

Hipel, K. W. and Mcleod, A. (1994) Time Series Modelling of Water Resources and Environmental Systems, Elsevier

**Examples**

```
data(tempEng)
```

# Index

\*Topic **datasets**

tempEng, 6

\*Topic **ts**

ardec, 1

ardec.lm, 3

ardec.periodic, 4

ardec.trend, 5

ar, 3

ardec, 1

ardec.lm, 3

ardec.periodic, 4

ardec.trend, 5

lm, 3

tempEng, 6