

Package ‘agriwater’

January 30, 2019

Title Evapotranspiration and Energy Fluxes Spatial Analysis

Version 1.0.0

Description Spatial modeling of energy balance and actual evapotranspiration using satellite images and meteorological data. Options of satellite are: Landsat-8 (with and without thermal bands), Sentinel-2 and MODIS. Respectively spatial resolutions are 30, 100, 10 and 250 meters. User can use data from a single meteorological station or a grid of meteorological stations (using any spatial interpolation method). Teixeira (2010) <doi:10.3390/rs0251287>. Teixeira et al. (2015) <doi:10.3390/rs71114597>. Silva, Manzione, and Albuquerque Filho (2018) <doi:10.3390/horticulturae4040044>.

Maintainer Cesar de Oliveira Ferreira Silva <cesaroliveira.f.silva@gmail.com>

Depends R (>= 3.2.0)

License MIT + file LICENSE

Encoding UTF-8

LazyData true

BugReports <https://github.com/cesarofs/agriwater/issues>

Imports raster, sp, rgdal

RoxygenNote 6.1.0

Suggests knitr, rmarkdown

VignetteBuilder knitr

NeedsCompilation no

Author Cesar de Oliveira Ferreira Silva [cre] (<<https://orcid.org/0000-0002-5152-6497>>), Antonio Heriberto de Castro Teixeira [aut], Rodrigo Lilla Manzione [ctb] (<<https://orcid.org/0000-0002-0754-2641>>)

Repository CRAN

Date/Publication 2019-01-30 22:34:00 UTC

R topics documented:

albedo_18	2
albedo_modis	3
albedo_s2	3
evapo_18	4
evapo_18t	4
evapo_18t_grid	5
evapo_18_grid	6
evapo_modis	6
evapo_modis_grid	8
evapo_s2	8
evapo_s2_grid	9
kc_18	9
kc_18t	10
kc_18t_grid	11
kc_18_grid	11
kc_modis	12
kc_modis_grid	13
kc_s2	14
kc_s2_grid	14
radiation_18	15
radiation_18t	15
radiation_18t_grid	16
radiation_18_grid	17
radiation_modis	17
radiation_modis_grid	19
radiation_s2	19
radiation_s2_grid	20
reflectance_18	21

Index

[22](#)

albedo_18

Surface Albedo using Landsat-8 images.

Description

Surface Albedo using Landsat-8 images.

Usage

`albedo_18(doy)`

Arguments

`doy` is the Day of Year (DOY)

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24").

albedo_modis

Surface Albedo using MODIS images.

Description

Surface Albedo using MODIS images.

Usage

albedo_modis()

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24").

albedo_s2

Surface Albedo using Sentinel-2 images.

Description

Surface Albedo using Sentinel-2 images.

Usage

albedo_s2()

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24").

evapo_18	<i>Actual evapotranspiration (ETa) using Landsat-8 images with single agrometeorological data.</i>
----------	--

Description

Actual evapotranspiration (ETa) using Landsat-8 images with single agrometeorological data.

Usage

```
evapo_18(doy, RG, Ta, ET0, a, b)
```

Arguments

doy	is the Day of Year (DOY)
RG	is the global solar radiation
Ta	is the average air temperature
ET0	is the reference evapotranspiration
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), net radiation ("Rn_MJ"), Crop Coefficient ("kc") and Actual Evapotranspiration (evapo).

evapo_18t	<i>Actual evapotranspiration (ETa) using Landsat-8 (including thermal bands) images with single agrometeorological data.</i>
-----------	--

Description

Actual evapotranspiration (ETa) using Landsat-8 (including thermal bands) images with single agrometeorological data.

Usage

```
evapo_18t(doy, RG, Ta, ET0, a, b)
```

Arguments

doy	is the Day of Year (DOY)
RG	is the global solar radiation
Ta	is the average air temperature
ET0	is the reference evapotranspiration
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), net radiation ("Rn_MJ"), Crop Coefficient ("kc") and Actual Evapotranspiration (evapo).

evapo_l8t_grid	<i>Actual evapotranspiration (ETa) using Landsat-8 (including thermal bands) images with a grid of agrometeorological data.</i>
----------------	---

Description

Actual evapotranspiration (ETa) using Landsat-8 (including thermal bands) images with a grid of agrometeorological data.

Usage

```
evapo_l8t_grid(doy, a, b)
```

Arguments

doy	is the Day of Year (DOY)
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), net radiation ("Rn_MJ"), Crop Coefficient ("kc") and Actual Evapotranspiration (evapo).

evapo_18_grid	<i>Actual evapotranspiration (ETa) using Landsat-8 images with a grid of agrometeorological data.</i>
---------------	---

Description

Actual evapotranspiration (ETa) using Landsat-8 images with a grid of agrometeorological data.

Usage

```
evapo_18_grid(doy, a, b)
```

Arguments

doy	is the Day of Year (DOY)
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), net radiation ("Rn_MJ"), Crop Coefficient ("kc") and Actual Evapotranspiration (evapo).

evapo_modis	<i>Actual evapotranspiration (ETa) using MODIS with single agrometeorological data.</i>
-------------	---

Description

Actual evapotranspiration (ETa) using MODIS with single agrometeorological data.

Usage

```
evapo_modis(doy, RG, Ta, ET0, a, b)
```

Arguments

doy	is the Day of Year (DOY)
RG	is the global solar radiation
Ta	is the average air temperature
ET0	is the reference evapotranspiration
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), net radiation ("Rn_MJ"), Crop Coefficient ("kc") and Actual Evapotranspiration (evapo).

Examples

```
library(agriwater)

# dependencies of package 'agriwater'
library(sp)
library(raster)
library(rgdal)

# Using a temporary folder to run example
wd <- tempdir()
initial = getwd()
setwd(wd)

# creating raster which simulate Sentinel-2 reflectances - for using
# real data, please download:
# https://drive.google.com/open?id=14E1wHNLxG7_Dh4I-GqNYakj8YJDgKLzk

wd <- tempdir()
xy <- matrix(rnorm(100, mean = 0.07, sd = 0.01), 10, 10)
rast <- raster(xy)
extent(rast) <- c(-40.5,-40.45,-9.5,-9.45)
projection(rast) <- CRS("+proj=longlat +datum=WGS84")
writeRaster(rast, file.path(wd, "B2"), format = "GTiff", overwrite=TRUE)
xy <- matrix(rnorm(100, mean = 0.05, sd = 0.015), 10, 10)
rast <- raster(xy)
extent(rast) <- c(-40.5,-40.45,-9.5,-9.45)
projection(rast) <- CRS("+proj=longlat +datum=WGS84")
writeRaster(rast, file.path(wd, "B1"), format = "GTiff", overwrite=TRUE)

mask <- as(extent(rast), 'SpatialPolygons')
projection(mask) <- CRS("+proj=longlat +datum=WGS84")
shapefile(mask, file.path(wd,"mask.shp"), overwrite=TRUE)

# using "agriwater" - it's the same procedure as the used for
# evapo_l8(), evapo_l8t(), evapo_modis_grid(), evapo_l8_grid(),
# evapo_l8t_grid(), evapo_s2() and evapo_s2_grid()
DOY = 134
a = 1.8
b = -0.008
RG = 17.6
Ta = 27.9
ET0 = 3.8
evapo_modis(DOY, RG, Ta, ET0, a, b)

#Exiting temporary folder and returning to previous workspace
```

```
setwd(initial)
```

<code>evapo_modis_grid</code>	<i>Actual evapotranspiration (ETa) using MODIS with a grid of agrometeorological data.</i>
-------------------------------	--

Description

Actual evapotranspiration (ETa) using MODIS with a grid of agrometeorological data.

Usage

```
evapo_modis_grid(doy, a, b)
```

Arguments

<code>doy</code>	is the Day of Year (DOY)
<code>a</code>	is one of the regression coefficients of SAFER algorithm
<code>b</code>	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), net radiation ("Rn_MJ"), Crop Coefficient ("kc") and Actual Evapotranspiration (evapo).

<code>evapo_s2</code>	<i>Actual evapotranspiration (ETa) using Sentinel-2 images with single agrometeorological data.</i>
-----------------------	---

Description

Actual evapotranspiration (ETa) using Sentinel-2 images with single agrometeorological data.

Usage

```
evapo_s2(doy, RG, Ta, ET0, a, b)
```

Arguments

<code>doy</code>	is the Day of Year (DOY)
<code>RG</code>	is the global solar radiation
<code>Ta</code>	is the average air temperature
<code>ET0</code>	is the reference evapotranspiration
<code>a</code>	is one of the regression coefficients of SAFER algorithm
<code>b</code>	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), net radiation ("Rn_MJ"), Crop Coefficient ("kc") and Actual Evapotranspiration (evapo).

evapo_s2_grid

Actual evapotranspiration (ETa) using Sentinel-2 images with a grid of agrometeorological data.

Description

Actual evapotranspiration (ETa) using Sentinel-2 images with a grid of agrometeorological data.

Usage

```
evapo_s2_grid(doy, a, b)
```

Arguments

- | | |
|-----|--|
| doy | is the Day of Year (DOY) |
| a | is one of the regression coefficients of SAFER algorithm |
| b | is one of the regression coefficients of SAFER algorithm |

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), net radiation ("Rn_MJ"), Crop Coefficient ("kc") and Actual Evapotranspiration (evapo).

kc_18

Crop coefficient (ETa / ET0) using Landsat-8 images with single agrometeorological data.

Description

Crop coefficient (ETa / ET0) using Landsat-8 images with single agrometeorological data.

Usage

```
kc_18(doy, RG, Ta, a, b)
```

Arguments

doy	is the Day of Year (DOY)
RG	is the global solar radiation
Ta	is the average air temperature
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc") and net radiation ("Rn_MJ").

kc_l8t

Crop coefficient (ETa / ET0) using Landsat-8 images (including thermal bands) with single agrometeorological data.

Description

Crop coefficient (ETa / ET0) using Landsat-8 images (including thermal bands) with single agrometeorological data.

Usage

```
kc_l8t(doy, RG, Ta, a, b)
```

Arguments

doy	is the Day of Year (DOY)
RG	is the global solar radiation
Ta	is the average air temperature
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc") and net radiation ("Rn_MJ").

kc_l8t_grid	<i>Crop coefficient (ETa / ET0) using Landsat-8 images (including thermal bands) with a grid of agrometeorological data.</i>
-------------	--

Description

Crop coefficient (ETa / ET0) using Landsat-8 images (including thermal bands) with a grid of agrometeorological data.

Usage

```
kc_l8t_grid(doy, a, b)
```

Arguments

- | | |
|-----|--|
| doy | is the Day of Year (DOY) |
| a | is one of the regression coefficients of SAFER algorithm |
| b | is one of the regression coefficients of SAFER algorithm |

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc") and net radiation ("Rn_MJ").

kc_l8_grid	<i>Crop coefficient (ETa / ET0) using Landsat-8 images with a grid of agrometeorological data.</i>
------------	--

Description

Crop coefficient (ETa / ET0) using Landsat-8 images with a grid of agrometeorological data.

Usage

```
kc_l8_grid(doy, a, b)
```

Arguments

- | | |
|-----|--|
| doy | is the Day of Year (DOY) |
| a | is one of the regression coefficients of SAFER algorithm |
| b | is one of the regression coefficients of SAFER algorithm |

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc") and net radiation ("Rn_MJ").

kc_modis

Crop coefficient (ETa / ET0) using MODIS with single agrometeorological data.

Description

Crop coefficient (ETa / ET0) using MODIS with single agrometeorological data.

Usage

```
kc_modis(doy, RG, Ta, a, b)
```

Arguments

doy	is the Day of Year (DOY)
RG	is the global solar radiation
Ta	is the average air temperature
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc") and net radiation ("Rn_MJ").

Examples

```
library(agriwater)

# dependencies of package 'agriwater'
library(sp)
library(raster)
library(rgdal)

# Using a temporary folder to run example
wd <- tempdir()
initial = getwd()
setwd(wd)

# creating raster which simulate MODIS reflectances - for using
# real data, please download:
# https://drive.google.com/open?id=14E1wHNLxG7_Dh4I-GqNYakj8YJDgKLzk

xy <- matrix(rnorm(100, mean = 0.07, sd = 0.01), 10, 10)
rast <- raster(xy)
extent(rast) <- c(-40.5,-40.45,-9.5,-9.45)
projection(rast) <- CRS("+proj=longlat +datum=WGS84")
```

```

writeRaster(rast, file.path(wd, "B2"), format = "GTiff", overwrite=TRUE)
xy <- matrix(rnorm(100, mean = 0.05, sd = 0.015), 10, 10)
rast <- raster(xy)
extent(rast) <- c(-40.5, -40.45, -9.5, -9.45)
projection(rast) <- CRS("+proj=longlat +datum=WGS84")
writeRaster(rast, file.path(wd, "B1"), format = "GTiff", overwrite=TRUE)

mask <- as(extent(rast), 'SpatialPolygons')
projection(mask) <- CRS("+proj=longlat +datum=WGS84")
shapefile(mask, file.path(wd, "mask.shp"), overwrite=TRUE)

# using "agriwater" - it's the same procedure as the used for
# kc_l8(), kc_l8t(), kc_s2(), kc_l8_grid(), kc_l8t_grid(),
# kc_s2(), kc_s2_grid() and kc_modis_grid()
DOY = 134
a = 1.8
b = -0.008
RG = 17.6
Ta = 27.9
kc_modis(DOY, RG, Ta, a, b)

#Exiting temporary folder and returning to previous workspace
setwd(initial)

```

kc_modis_grid

Crop coefficient (ET_a / ET₀) using MODIS with a grid of agrometeorological data.

Description

Crop coefficient (ET_a / ET₀) using MODIS with a grid of agrometeorological data.

Usage

```
kc_modis_grid(doy, a, b)
```

Arguments

- | | |
|-----|--|
| doy | is the Day of Year (DOY) |
| a | is one of the regression coefficients of SAFER algorithm |
| b | is one of the regression coefficients of SAFER algorithm |

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc") and net radiation ("Rn_MJ").

<code>kc_s2</code>	<i>Crop coefficient (ETa / ET0) using Sentinel-2 images with single agrometeorological data.</i>
--------------------	--

Description

Crop coefficient (ETa / ET0) using Sentinel-2 images with single agrometeorological data.

Usage

```
kc_s2(doy, RG, Ta, a, b)
```

Arguments

doy	is the Day of Year (DOY)
RG	is the global solar radiation
Ta	is the average air temperature
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc") and net radiation ("Rn_MJ").

<code>kc_s2_grid</code>	<i>Crop coefficient (ETa / ET0) using Sentinel-2 images with a grid of agrometeorological data.</i>
-------------------------	---

Description

Crop coefficient (ETa / ET0) using Sentinel-2 images with a grid of agrometeorological data.

Usage

```
kc_s2_grid(doy, a, b)
```

Arguments

doy	is the Day of Year (DOY)
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc") and net radiation ("Rn_MJ").

radiation_18	<i>Energy balance using Landsat-8 images with single agrometeorological data.</i>
--------------	---

Description

Energy balance using Landsat-8 images with single agrometeorological data.

Usage

```
radiation_18(doy, RG, Ta, ET0, a, b)
```

Arguments

doy	is the Day of Year (DOY)
RG	is the global solar radiation
Ta	is the average air temperature
ET0	is the reference evapotranspiration
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc"), Actual Evapotranspiration (evapo), latent heat flux "LE_MJ", net radiation ("Rn_MJ"), ground heat flux ("G_MJ") and the sensible heat flux ("H_MJ").

radiation_18t	<i>Energy balance using Landsat-8 images (including thermal bands) with single agrometeorological data.</i>
---------------	---

Description

Energy balance using Landsat-8 images (including thermal bands) with single agrometeorological data.

Usage

```
radiation_18t(doy, RG, Ta, ET0, a, b)
```

Arguments

doy	is the Day of Year (DOY)
RG	is the global solar radiation
Ta	is the average air temperature
ET0	is the reference evapotranspiration
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc"), Actual Evapotranspiration (evapo), latent heat flux "LE_MJ", net radiation ("Rn_MJ"), ground heat flux ("G_MJ") and the sensible heat flux ("H_MJ").

radiation_l8t_grid *Energy balance using Landsat-8 images (including thermal bands) with a grid of agrometeorological data.*

Description

Energy balance using Landsat-8 images (including thermal bands) with a grid of agrometeorological data.

Usage

```
radiation_l8t_grid(doy, a, b)
```

Arguments

doy	is the Day of Year (DOY)
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc"), Actual Evapotranspiration (evapo), latent heat flux "LE_MJ", net radiation ("Rn_MJ"), ground heat flux ("G_MJ") and the sensible heat flux ("H_MJ").

radiation_l8_grid	<i>Energy balance using Landsat-8 images with a grid of agrometeorological data.</i>
-------------------	--

Description

Energy balance using Landsat-8 images with a grid of agrometeorological data.

Usage

```
radiation_l8_grid(doy, a, b)
```

Arguments

doy	is the Day of Year (DOY)
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc"), Actual Evapotranspiration (evapo), latent heat flux "LE_MJ", net radiation ("Rn_MJ"), ground heat flux ("G_MJ") and the sensible heat flux ("H_MJ").

radiation_modis	<i>Energy balance using Landsat-8 images with single agrometeorological data.</i>
-----------------	---

Description

Energy balance using Landsat-8 images with single agrometeorological data.

Usage

```
radiation_modis(doy, RG, Ta, ET0, a, b)
```

Arguments

doy	is the Day of Year (DOY)
RG	is the global solar radiation
Ta	is the average air temperature
ET0	is the reference evapotranspiration
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc"), Actual Evapotranspiration (evapo), latent heat flux "LE_MJ", net radiation ("Rn_MJ"), ground heat flux ("G_MJ") and the sensible heat flux ("H_MJ").

Examples

```
library(agriwater)

# dependencies of package 'agriwater'
library(sp)
library(raster)
library(rgdal)

# Using a temporary folder to run example
wd <- tempdir()
initial = getwd()
setwd(wd)

# creating raster which simulate Sentinel-2 reflectances - for using
# real data, please download:
# https://drive.google.com/open?id=14E1wHNLxG7\_Dh4I-GqNYakj8YJDgKLzk

wd <- tempdir()
setwd <- tempdir()
xy <- matrix(rnorm(100, mean = 0.07, sd = 0.01), 10, 10)
rast <- raster(xy)
extent(rast) <- c(-40.5,-40.45,-9.5,-9.45)
projection(rast) <- CRS("+proj=longlat +datum=WGS84")
writeRaster(rast, file.path(wd, "B2"), format = "GTiff", overwrite=TRUE)
xy <- matrix(rnorm(100, mean = 0.05, sd = 0.015), 10, 10)
rast <- raster(xy)
extent(rast) <- c(-40.5,-40.45,-9.5,-9.45)
projection(rast) <- CRS("+proj=longlat +datum=WGS84")
writeRaster(rast, file.path(wd, "B1"), format = "GTiff", overwrite=TRUE)

# creating mask of study area
mask <- as(extent(rast), 'SpatialPolygons')
projection(mask) <- CRS("+proj=longlat +datum=WGS84")
shapefile(mask, file.path(wd,"mask.shp"), overwrite=TRUE)

# using "agriwater" - it's the same procedure as the used for
# radiation_l8(), radiation_l8t(), radiation_s2(),
# radiation_l8_grid(), radiation_l8t_grid(),
# radiation_s2_grid(), radiation_s2() and radiation_modis_grid()
DOY = 134
a = 1.8
b = -0.008
RG = 17.6
Ta = 27.9
ET0 = 3.8
radiation_modis(DOY, RG, Ta, ET0, a, b)
```

```
#Exiting temporary folder and returning to previous workspace  
setwd(initial)
```

radiation_modis_grid *Energy balance using Landsat-8 images with a grid of agrometeorological data.*

Description

Energy balance using Landsat-8 images with a grid of agrometeorological data.

Usage

```
radiation_modis_grid(doy, a, b)
```

Arguments

- | | |
|-----|--|
| doy | is the Day of Year (DOY) |
| a | is one of the regression coefficients of SAFER algorithm |
| b | is one of the regression coefficients of SAFER algorithm |

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc"), Actual Evapotranspiration (evapo), latent heat flux "LE_MJ", net radiation ("Rn_MJ"), ground heat flux ("G_MJ") and the sensible heat flux ("H_MJ").

radiation_s2 *Energy balance using Sentinel-2 images with single agrometeorological data.*

Description

Energy balance using Sentinel-2 images with single agrometeorological data.

Usage

```
radiation_s2(doy, RG, Ta, ET0, a, b)
```

Arguments

doy	is the Day of Year (DOY)
RG	is the global solar radiation
Ta	is the average air temperature
ET0	is the reference evapotranspiration
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc"), Actual Evapotranspiration (evapo), latent heat flux "LE_MJ", net radiation ("Rn_MJ"), ground heat flux ("G_MJ") and the sensible heat flux ("H_MJ").

radiation_s2_grid	<i>Energy balance using Sentinel-2 images with a grid of agrometeorological data.</i>
--------------------------	---

Description

Energy balance using Sentinel-2 images with a grid of agrometeorological data.

Usage

```
radiation_s2_grid(doy, a, b)
```

Arguments

doy	is the Day of Year (DOY)
a	is one of the regression coefficients of SAFER algorithm
b	is one of the regression coefficients of SAFER algorithm

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24"), NDVI, Surface Temperature ("LST"), Crop Coefficient ("kc"), Actual Evapotranspiration (evapo), latent heat flux "LE_MJ", net radiation ("Rn_MJ"), ground heat flux ("G_MJ") and the sensible heat flux ("H_MJ").

reflectance_l8 *Reflectancies from Landsat-8 images.*

Description

Reflectancies from Landsat-8 images.

Usage

`reflectance_l8(doy)`

Arguments

doy is the Day of Year (DOY)

Value

It returns in raster format (.tif) the Surface Albedo at 24h scale ("Alb_24").

Index

albedo_l8, 2
albedo_modis, 3
albedo_s2, 3

evapo_l8, 4
evapo_l8_grid, 6
evapo_l8t, 4
evapo_l8t_grid, 5
evapo_modis, 6
evapo_modis_grid, 8
evapo_s2, 8
evapo_s2_grid, 9

kc_l8, 9
kc_l8_grid, 11
kc_l8t, 10
kc_l8t_grid, 11
kc_modis, 12
kc_modis_grid, 13
kc_s2, 14
kc_s2_grid, 14

radiation_l8, 15
radiation_l8_grid, 17
radiation_l8t, 15
radiation_l8t_grid, 16
radiation_modis, 17
radiation_modis_grid, 19
radiation_s2, 19
radiation_s2_grid, 20
reflectance_l8, 21