

A tutorial dbEmpLikeGOF R package

Jeffrey C. Miecznikowski*, Albert Vexler†, and Lori A. Shepherd◇

August 23, 2013

Statistical Genetics and Genomics Research Group
Department of Biostatistics, University at Buffalo
New York State Center of Excellence in Bioinformatics and Life Sciences
Roswell Park Cancer Institute

*jcm38@buffalo.edu, †avexler@buffalo.edu, ◇las65@buffalo.edu

Contents

1	Introduction	2
2	Examples	2
3	References	4

1 Introduction

The dbEmplikeGOF package provides a function dbEmplikeGOF to be used for density based empirical likelihood (EL) goodness-of-fit tests based on sample entropy, as well as to perform the two sample EL ratio test for distribution equality. The function provides the test statistic and associated p-values. The p-value can be calculated by Monte-Carlo methods or estimated based on pre-calculated tables of selected sample sizes and alpha values. For details and algorithms:

Jeffrey C. Miecznikowski, Albert Vexler, Lori A. Shepherd (2013). dbEmpLikeGOF: An R Package for Nonparametric Likelihood Ratio Tests for Goodness-of-Fit and Two-Sample Comparisons Based on Sample Entropy. *Journal of Statistical Software*, 54(3), 1-19.

Vexler A, Gurevich G, Empirical likelihood ratios applied to goodness-of-fit tests based on sample entropy. *Computational Statistics and Data Analysis* 54(2010) 531-545.

Gurevich G, Vexler A, A two-sample empirical likelihood ratio test based on sample entropy. *Statistics and Computing*, 2011.

2 Examples

The following performs a density-based empirical likelihood based goodness-of-fit tests based on sample entropy and calculates the p-value based on Monte-Carlo methods. The examples examine three null hypothesis, 1) data follows a normal distribution with unknown mean and standard deviation, 2) data follows a uniform distribution on 0 to 1 and 3) data from two samples are from the same distribution. The example below tests the data (`normData`) against the normal distribution.

```
> library(dbEmpLikeGOF)
> normData = rnorm(25)
> dbEmpLikeGOF(x=normData, testcall="normal", pvl.Table=FALSE)

...Working on teststat
...Working on p-value
$teststat
[1] 3.746553

$pvalue
[1] 0.981

>
```

The p-value can be estimated based on precalculated tables rather than performing Monte-Carlo methods. This is controlled by the argument `pvl.Table`. To estimate based on tables `pvl.Table` argument is `TRUE`, which is the default setting.

```
> dbEmpLikeGOF(x=normData, testcall="normal", pvl.Table=TRUE)

...Working on teststat
estimating pvalue based on table
$teststat
[1] 3.746553

$pvalue
[1] 0.9848558

>
```

Similar calculations can be made to test data against a uniform distribution on zero to one.

```
> unifData = runif(30)
> # calculates pvalue based on Monte-Carlo methods
> dbEmpLikeGOF(x=unifData, testcall="uniform", pvl.Table=FALSE)

...Working on teststat
...Working on p-value
$teststat
[1] 8.692456

$pvalue
[1] 0.064

> # estimates pvalue based on tables
> dbEmpLikeGOF(x=unifData, testcall="uniform", pvl.Table=TRUE)

...Working on teststat
estimating pvalue based on table
$teststat
[1] 8.692456

$pvalue
[1] 0.06285791

>
```

Notice the data in each of the above examples was designed to match the proposed distribution. Below is an example where the data does not follow the proposed distribution

```

> dbEmpLikeGOF(x=unifData, testcall="normal", pv1.Table=TRUE)
...Working on teststat
estimating pvalue based on table
$teststat
[1] 11.79232

$pvalue
[1] 0.005150142
>

```

It is also possible to test for distribution equality between two samples. When specifying an x and y samples, the dbEmpLikeGOF function will test for distribution equality between the two samples.

```

> dbEmpLikeGOF(x=unifData, y=normData, pv1.Table=TRUE)
...Working on teststat
estimating pvalue based on table
$teststat
[1] 23.42204

$pvalue
[1] 0.001

> normDataSet2 = rnorm(40)
> dbEmpLikeGOF(x=normDataSet2, y=normData, pv1.Table=TRUE)
...Working on teststat
estimating pvalue based on table
$teststat
[1] 10.94152

$pvalue
[1] 0.4913884
>

```

Notice the sample vectors do not have to be of equal length.

3 References

For additional details and examples please see:

Jeffrey C. Miecznikowski, Albert Vexler, Lori A. Shepherd (2013). dbEmpLikeGOF: An R Package for Nonparametric Likelihood Ratio Tests for Goodness-of-Fit and Two-Sample Comparisons Based on Sample Entropy. *Journal of Statistical Software*, 54(3), 1-19.

<http://www.jstatsoft.org/v54/i03/>