

# Package ‘rlfsm’

November 5, 2019

**Type** Package

**Title** Simulations and Statistical Inference for Linear Fractional Stable Motions

**Version** 1.0.0

**Maintainer** Dmitry Otryakhin <d.otryakhin.acad@protonmail.ch>

**Author** Dmitry Otryakhin [aut, cre],  
Stepan Mazur [aut],  
Mathias Ljungdahl [ctb]

**Description**

Contains functions for simulating linear fractional stable motions, according to techniques developed by Stoev and Taqqu (2004) <doi:10.1142/S0218348X04002379>, as well as functions for computing important statistics used with these processes introduced by Mazur, Otryakhin and Podolskij (2018) <arXiv:1802.06373>, and also different quantities related to those statistics.

**License** GPL-3

**URL** [https://gitlab.com/Dmitry\\_Otryakhin/  
Tools-for-parameter-estimation-of-the-linear-fractional-stable-motion](https://gitlab.com/Dmitry_Otryakhin/Tools-for-parameter-estimation-of-the-linear-fractional-stable-motion)

**Encoding** UTF-8

**LazyData** true

**RoxygenNote** 6.1.1

**Depends** methods, foreach, doParallel

**Imports** ggplot2, stabledist, reshape2, plyr, Rdpack

**Suggests** elliptic, spatstat, testthat, stringi

**RdMacros** Rdpack

**NeedsCompilation** no

**Repository** CRAN

**Date/Publication** 2019-11-05 13:40:02 UTC

## R topics documented:

alpha_hat . . . . .	2
a_p . . . . .	4
a_tilda . . . . .	4
CLT . . . . .	5
ContinEstim . . . . .	6
GenHighEstim . . . . .	7
GenLowEstim . . . . .	8
H_hat . . . . .	9
h_kr . . . . .	9
increment . . . . .	10
MCestimLFSM . . . . .	11
MinContrastEstim . . . . .	13
m_pk . . . . .	14
Norm_alpha . . . . .	15
path . . . . .	16
paths . . . . .	17
Path_array . . . . .	18
phi . . . . .	19
phi_of_alpha . . . . .	20
Plot_dens . . . . .	20
Plot_list_paths . . . . .	21
Plot_vb . . . . .	22
Retrieve_stats . . . . .	23
R_hl . . . . .	24
sf . . . . .	25
sigma_hat . . . . .	26
theta . . . . .	27
U_g . . . . .	28
U_gh . . . . .	29
U_ghuv . . . . .	29
	31

## Index

---

alpha_hat	<i>Statistical estimator for alpha</i>
-----------	--

---

### Description

Defined for the two frequencies as

$$\widehat{\alpha}_{high} := \frac{\log |\log \varphi_{high}(t_2; \widehat{H}_{high}(p, k)_n, k)_n| - \log |\log \varphi_{high}(t_1; \widehat{H}_{high}(p, k)_n, k)_n|}{\log t_2 - \log t_1}$$

$$\widehat{\alpha}_{low} := \frac{\log |\log \varphi_{low}(t_2; k)_n| - \log |\log \varphi_{low}(t_1; k)_n|}{\log t_2 - \log t_1}$$

**Usage**

```
alpha_hat(t1, t2, k, path, H, freq)
```

**Arguments**

t1	real number such that $t_2 > t_1 > 0$
t2	real number such that $t_2 > t_1 > 0$
k	increment order
path	sample path of l fsm on which the inference is to be performed
H	Hurst parameter
freq	Frequency of the motion. It can take two values: "H" for high frequency and "L" for the low frequency setting.

**Details**

The function triggers function [phi](#), thus Hurst parameter is required only in high frequency case. In the low frequency, there is no need to assign H a value because it will not be evaluated.

**References**

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

**Examples**

```
m<-45; M<-60; N<-2^14-M
alpha<-1.8; H<-0.8; sigma<-0.3
freq='H'
r=1; k=2; p=0.4; t1=1; t2=2

# Estimating alpha in the high frequency case
# using preliminary estimation of H
l fsm<-path(N=N,m=m,M=M,alpha=alpha,H=H,
            sigma=sigma,freq='L',disable_X=FALSE,seed=3)$l fsm

H_est<-H_hat(p=p,k=k,path=l fsm)
H_est
alpha_est<-alpha_hat(t1=t1,t2=t2,k=k,path=l fsm,H=H_est,freq=freq)
alpha_est
```

<code>a_p</code>	<i>Function a_p.</i>
------------------	----------------------

### Description

Computes the corresponding function value from Mazur et al. 2018.

### Usage

`a_p(p)`

### Arguments

<code>p</code>	power, real number from (-1,1)
----------------	--------------------------------

### References

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

<code>a_tilda</code>	<i>Creates the corresponding value from the paper by Stoev and Taqqu (2004).</i>
----------------------	--

### Description

Creates the corresponding value from the paper by Stoev and Taqqu (2004).

### Usage

`a_tilda(N, m, M, alpha, H)`

### Arguments

<code>N</code>	a number of points of the l fsm.
<code>m</code>	discretization. A number of points between two nearby motion points
<code>M</code>	truncation parameter. A number of points at which the integral representing the definition of l fsm is calculated. So, after M points back we consider the rest of the integral to be 0.
<code>alpha</code>	self-similarity parameter of alpha stable random motion.
<code>H</code>	Hurst parameter

### References

Stoev S, Taqqu M (2004). “Simulation methods for linear fractional stable motion and FARIMA using the Fast Fourier Transform.” *Fractals*, **95**(1), 95-121.

---

CLT	<i>The function explores numerical properties of statistical estimators operating on random processes. Deprecated.</i>
-----	--

---

## Description

The function is left for backward compatibility. The newer version of it is [MCestimLFSM](#). The function is useful, for instance, when one needs to compute standard deviation of  $\hat{\alpha}_{high}$  estimator given a fixed set of parameters.

## Usage

```
CLT(Nmc, s, m, M, alpha, H, sigma, fr, Inference, ...)
```

## Arguments

Nmc	Number of Monte Carlo repetitions
s	sequence of path lengths
m	discretization. A number of points between two nearby motion points
M	truncation parameter. A number of points at which the integral representing the definition of l fsm is calculated. So, after M points back we consider the rest of the integral to be 0.
alpha	self-similarity parameter of alpha stable random motion.
H	Hurst parameter
sigma	Scale parameter of l fsm
fr	frequency. Either "H" or "L"
Inference	statistical function to apply to sample paths
...	parameters to pass to Inference

## Details

CLT performs Monte-Carlo experiments to compute parameters according to procedure Inference. More specifically, for each element of s it generates Nmc l fsm sample paths with length equal to s[i], performs the statistical inference on each, obtaining the estimates, and then returns their different statistics. It is vital that the estimator returns a list of named parameters (one or several of 'sigma', 'alpha' and 'H'). CLT uses the names to lookup the true parameter value and compute its bias.

For sample path generation CLT uses a light-weight version of path, path\_fast. In order to be applied, function Inference must accept argument 'path' as a sample path.

## Value

It returns a list containing the following components:

CLT_dataset	a data frame, standardized values of the estimates depending on path length s
BSdM	a data frame, means, biases and standard deviations depending on s
Inference	a closure used to obtain estimates
alpha, H, sigma	the parameters for which CLT performs path generation
freq	frequency, either 'L' for low- or 'H' for high frequency

ContinEstim

*Parameter estimation procedure in continuous case.*

## Description

Parameter freq is preserved to allow for investigation of the inference procedure in high frequency case.

## Usage

```
ContinEstim(t1, t2, p, k, path, freq)
```

## Arguments

t1, t2	real number such that $t2 > t1 > 0$
p	power
k	increment order
path	sample path of l fsm on which the inference is to be performed
freq	Frequency of the motion. It can take two values: "H" for high frequency and "L" for the low frequency setting.

## References

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

## Examples

```
m<-45; M<-60; N<-2^10-M
alpha<-0.8; H<-0.8; sigma<-0.3
p<-0.3; k=3; t1=1; t2=2

l fsm<-path(N=N,m=m,M=M,alpha=alpha,H=H,
            sigma=sigma,freq='L',disable_X=FALSE,seed=3)$l fsm
ContinEstim(t1,t2,p,k,path=l fsm,freq='L')
```

---

GenHighEstim*High frequency estimation procedure for l fsm.*

---

## Description

General estimation procedure for high frequency case when 1/alpha is not a natural number. "Unnecessary" parameter freq is preserved to allow for investigation of the inference procedure in low frequency case

## Usage

```
GenHighEstim(p, p_prime, path, freq, low_bound = 0.01, up_bound = 4)
```

## Arguments

p	power
p_prime	power
path	sample path of l fsm on which the inference is to be performed
freq	Frequency of the motion. It can take two values: "H" for high frequency and "L" for the low frequency setting.
low_bound	positive real number
up_bound	positive real number

## Details

In this algorithm the preliminary estimate of alpha is found via using [uniroot](#) function. The latter is given the lower and the upper bounds for alpha via low\_bound and up\_bound parameters. It is not possible to pass 0 as the lower bound because there are numerical limitations on the alpha estimate, caused by the length of the sample path and by numerical errors. p and p\_prime must belong to the interval (0,1/2) (in the notation kept in rlfsm package) The two powers cannot be equal.

## References

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

## Examples

```
m<-45; M<-60; N<-2^10-M
sigma<-0.3
p<-0.2; p_prime<-0.4

#### Continuous case
l fsm<-path(N=N,m=m,M=M,alpha=1.8,H=0.8,
           sigma=sigma,freq='L',disable_X=FALSE,seed=3)$l fsm

GenHighEstim(p=p,p_prime=p_prime,path=l fsm,freq="H")
```

```
#### H-1/alpha<0 case
lfsm<-path(N=N, m=m, M=M, alpha=0.8, H=0.8,
            sigma=sigma, freq='H', disable_X=FALSE, seed=3)$lfsm

GenHighEstim(p=p, p_prime=p_prime, path=lfsm, freq="H")
```

**GenLowEstim***Low frequency estimation procedure for lfsm.***Description**

General estimation procedure for low frequency case when  $1/\alpha$  is not a natural number.

**Usage**

```
GenLowEstim(t1, t2, p, path, freq = "L")
```

**Arguments**

t1	real number such that $t_2 > t_1 > 0$
t2	real number such that $t_2 > t_1 > 0$
p	power
path	sample path of lfsm on which the inference is to be performed
freq	Frequency of the motion. It can take two values: "H" for high frequency and "L" for the low frequency setting.

**References**

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

**Examples**

```
m<-45; M<-60; N<-2^10-M
sigma<-0.3
p<-0.3; k=3; t1=1; t2=2

#### Continuous case
lfsm<-path(N=N, m=m, M=M, alpha=1.8, H=0.8,
            sigma=sigma, freq='L', disable_X=FALSE, seed=3)$lfsm

GenLowEstim(t1=t1, t2=t2, p=p, path=lfsm, freq="L")

#### H-1/alpha<0 case
lfsm<-path(N=N, m=m, M=M, alpha=0.8, H=0.8,
            sigma=sigma, freq='L', disable_X=FALSE, seed=3)$lfsm
```

---

```
GenLowEstim(t1=t1,t2=t2,p=p,path=lfsm,freq="L")

##### The procedure works also for high frequency case
lfsm<-path(N=N,m=m,M=M,alpha=1.8,H=0.8,
              sigma=sigma,freq='H',disable_X=FALSE,seed=3)$lfsm

GenLowEstim(t1=t1,t2=t2,p=p,path=lfsm,freq="H")
```

---

**H\_hat***Statistical estimator of H in high/low frequency setting***Description**

The statistic is defined as

$$\hat{H}_{\text{high}}(p, k)_n := \frac{1}{p} \log_2 R_{\text{high}}(p, k)_n, \quad \hat{H}_{\text{low}}(p, k)_n := \frac{1}{p} \log_2 R_{\text{low}}(p, k)_n$$

**Usage**

```
H_hat(p, k, path)
```

**Arguments**

p	power
k	increment order
path	sample path of lfsm on which the inference is to be performed

**References**

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

---

**h\_kr***Function h\_kr*

---

**Description**

Function  $h_{k,r} : R \rightarrow R$  is given by

$$h_{k,r}(x) = \sum_{j=0}^k (-1)^j \binom{k}{j} (x - rj)_+^{H-1/\alpha}, \quad x \in R$$

**Usage**

```
h_kr(k, r, x, H, alpha, l = 0)
```

## Arguments

k	order of the increment, a natural number
r	difference step, a natural number
x	real number
H	Hurst parameter
alpha	self-similarity parameter of alpha stable random motion.
l	a value by which we shift x. Is used for computing function f_.+l and is passed to integrate function.

## References

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

## Examples

```
#### Plot h_kr ####
s<-seq(0,10, by=0.01)
h_val<-sapply(s,h_kr, k=5, r=1, H=0.3, alpha=1)
plot(s,h_val)
```

increment	<i>Higher order increments</i>
-----------	--------------------------------

## Description

Difference of the kth order. Defined as following:

$$\Delta_{i,k}^{n,r} X := \sum_{j=0}^k (-1)^j \binom{k}{j} X_{(i-rj)/n}, i \geq rk.$$

Index i here is a coordinate in terms of point\_num. Although R uses vector indexes that start from 1, increment has i varying from 0 to N, so that a vector has a length N+1. It is done in order to comply with the notation of the paper. This function doesn't allow for choosing frequency n. The frequency is determined by the path supplied, thus n equals to either the length of the path in high frequency setting or 1 in low frequency setting. increment() gives increments at certain point passed as i, which is a vector here. increments() computes high order increments for the whole sample path. The first function evaluates the formula above, while the second one uses structure diff(diff(...)) because the formula is slower at higher k.

## Usage

```
increment(r, i, k, path)

increments(k, r, path)
```

## Arguments

r	difference step, a natural number
i	index of the point at which the increment is to be computed, a natural number.
k	order of the increment, a natural number
path	sample path for which a kth order increment is computed

## References

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

## Examples

```
m<-45; M<-60; N<-2^10-M
alpha<-0.8; H<-0.8; sigma<-0.3

l fsm<-path(N=N,m=m,M=M,alpha=alpha,H=H,
             sigma=sigma,freq='L',disable_X=FALSE,seed=3)$l fsm

tryCatch(
  increment(r=1,i=length(l fsm),k=length(l fsm)+100,path=l fsm),
  error=function(c) 'An error occurs when k is larger than the length of the sample path')

increment(r=3,i=50,k=3,path=l fsm)

path=c(1,4,3,6,8,5,3,5,8,5,1,8,6)

r=2; k=3
n <- length(path) - 1
DeltaX = increment(seq(r*k, n), path = path, k = k, r = r)
DeltaX == increments(k=k,r=r,path)
```

## Description

The function is useful, for instance, when one needs to compute standard deviation of  $\hat{\alpha}_{high}$  estimator given a fixed set of parameters.

## Usage

```
MCestimLFSM(Nmc, s, m, M, alpha, H, sigma, fr, Inference, ...)
```

## Arguments

Nmc	Number of Monte Carlo repetitions
s	sequence of path lengths
m	discretization. A number of points between two nearby motion points
M	truncation parameter. A number of points at which the integral representing the definition of l fsm is calculated. So, after M points back we consider the rest of the integral to be 0.
alpha	self-similarity parameter of alpha stable random motion.
H	Hurst parameter
sigma	Scale parameter of l fsm
fr	frequency. Either "H" or "L"
Inference	statistical function to apply to sample paths
...	parameters to pass to Inference

## Details

MCestimLFSM performs Monte-Carlo experiments to compute parameters according to procedure Inference. More specifically, for each element of s it generates Nmc l fsm sample paths with length equal to s[i], performs the statistical inference on each, obtaining the estimates, and then returns their different statistics. It is vital that the estimator returns a list of named parameters (one or several of 'sigma', 'alpha' and 'H'). MCestimLFSM uses the names to lookup the true parameter value and compute its bias.

For sample path generation MCestimLFSM uses a light-weight version of path, path\_fast. In order to be applied, function Inference must accept argument 'path' as a sample path.

## Value

It returns a list containing the following components:

data	a data frame, values of the estimates depending on path length s
data_nor	a data frame, normalized values of the estimates depending on path length s
means, biases, sds	data frames: means, biases and standard deviations of the estimators depending on s
Inference	a function used to obtain estimates
alpha, H, sigma	the parameters for which MCestimLFSM performs path generation
freq	frequency, either 'L' for low- or 'H' for high frequency

## Examples

```
#### Set of global parameters ####
m<-25; M<-60
p<-.4; p_prime<-.2; k<-2
t1<-1; t2<-2
NmonteC<-5e1
```

```

S<-c(1e2,3e2)
alpha<-1.8; H<-0.8; sigma<-0.3

# How to plot empirical density

theor_3_1_H_clt<-MCestimLFSM(s=S,fr='H',Nmc=NmonteC,
                                m=m,M=M,alpha=alpha,H=H,
                                sigma=sigma,ContinEstim,
                                t1=t1,t2=t2,p=p,k=k)
l_plot<-Plot_dens(par_vec=c('sigma','alpha','H'),
                    MC_data=theor_3_1_H_clt, Nnorm=1e7)

# For MCestimLFSM() it is vital that the estimator returns a list of named parameters

H_hat_f <- function(p,k,path) {hh<-H_hat(p,k,path); list(H=hh)}
theor_3_1_H_clt<-MCestimLFSM(s=S,fr='H',Nmc=NmonteC,
                                m=m,M=M,alpha=alpha,H=H,
                                sigma=sigma,H_hat_f,
                                p=p,k=k)

# The estimator can return one, two or three of the parameters.

est_1 <- function(path) list(H=1)
theor_3_1_H_clt<-MCestimLFSM(s=S,fr='H',Nmc=NmonteC,
                                m=m,M=M,alpha=alpha,H=H,
                                sigma=sigma,est_1)

est_2 <- function(path) list(H=0.8, alpha=1.5)
theor_3_1_H_clt<-MCestimLFSM(s=S,fr='H',Nmc=NmonteC,
                                m=m,M=M,alpha=alpha,H=H,
                                sigma=sigma,est_2)

est_3 <- function(path) list(sigma=5, H=0.8, alpha=1.5)
theor_3_1_H_clt<-MCestimLFSM(s=S,fr='H',Nmc=NmonteC,
                                m=m,M=M,alpha=alpha,H=H,
                                sigma=sigma,est_3)

```

**MinContrastEstim**

*Statistical estimator of sigma, alpha and H in low frequency setting based on minimal contrast estimation comparing the empirical characteristic function with the true one*

**Description**

Estimates H using the [H\\_hat](#) function while sigma and alpha are obtained via

$$\arg \min_{\sigma, \alpha} \int_0^{\infty} (\varphi_n(t) - \varphi_{\sigma, \alpha, H_{hat}}(t))^2 \exp(-t^2/2) dt$$

, where  $\varphi_n$  is the empirical characteristic function, see [phi](#), and  $\varphi_{\sigma,\alpha,H_{hat}}$  is the characteristic function of the kth order increment wrt the parameters  $\sigma, \alpha, H_{hat}$ , see also [increment](#).

### Usage

```
MinContrastEstim(path, k, p, order_GH)
```

### Arguments

path	low frequency sample path from which the parameters should be estimated.
k	order of increments.
p	any real number, the power used for <a href="#">H_hat</a> .
order_GH	number of weights in the Gauss-Hermite approximation of the integral, see the <a href="#">gauss.hermite</a> function from the <a href="#">spatstat</a> package.

### Details

This algorithm approximates the above integral using Gauss-Hermite quadrature and uses the L-BFGS-B method from the [optim](#) function to minimize over the parameters sigma and alpha. Due to numerical problems estimation of sigma below 0.01 and alpha or H below 0.05 is currently not possible.

### References

Ljungdahl MM, Podolskij M (2019). “A Minimal Contrast Estimator for the Linear Fractional Stable Motion.” *To be submitted*.

### Examples

```
m0 = 256
M0 = 600
alpha0 = 1.8
H0 = 0.8
sigma0 = 0.3
n = 100
X <- path(N = n, m = m0, M = M0, alpha = alpha0, H = H0, sigma = sigma0, freq = 'L')$lfsm
MinContrastEstim(path = X, k = 2, p = 0.4, order_GH = 8)
```

*m\_pk*

$m(-p,k)$

### Description

defined as  $m_{p,k} := E[|\Delta_{k,k} X|^p]$  for positive powers. When p is negative (-p is positive) the equality does not hold.

### Usage

```
m_pk(k, p, alpha, H, sigma)
```

**Arguments**

k	increment order
p	a positive number
alpha	self-similarity parameter of alpha stable random motion.
H	Hurst parameter
sigma	Scale parameter of Ifsm

**Details**

The following identity is used for computations:

$$m_{-p,k} = \frac{(\sigma \|h_k\|_\alpha)^{-p}}{a_{-p}} \int_{\mathbb{R}} \exp(-|y|^\alpha) |y|^{-1+p} dy = \frac{2(\sigma \|h_k\|_\alpha)^{-p}}{\alpha a_{-p}} \Gamma(p/\alpha)$$

**References**

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

Norm\_alpha

*Alpha-norm of an arbitrary function***Description**

Alpha-norm of an arbitrary function

**Usage**

```
Norm_alpha(fun, alpha, ...)
```

**Arguments**

fun	a function to compute a norm
alpha	self-similarity parameter of alpha stable random motion.
...	a set of parameters to pass to integrate

**Details**

fun must accept a vector of values for evaluation. See ?integrate for further details. Most problems with this function appear because of rather high precision. Try to tune rel.tol parameter first.

**References**

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

**Examples**

```
Norm_alpha(h_kr, alpha=1.8, k=2, r=1, H=0.8, l=4)
```

---

path	<i>Generator of linear fractional stable motion</i>
------	---

---

### Description

The function creates a 1-dimensional l fsm sample path using the numerical algorithm from the paper by Stoev and Taqqu. Linear fractional stable motion is defined as

$$X_t = \int_{\mathbb{R}} \left\{ (t-s)_+^{H-1/\alpha} - (-s)_+^{H-1/\alpha} \right\} dL_s$$

### Usage

```
path(N = NULL, m, M, alpha, H, sigma, freq, disable_X = FALSE,
      levy_increments = NULL, seed = NULL)
```

### Arguments

N	a number of points of the l fsm.
m	discretization. A number of points between two nearby motion points
M	truncation parameter. A number of points at which the integral representing the definition of l fsm is calculated. So, after M points back we consider the rest of the integral to be 0.
alpha	self-similarity parameter of alpha stable random motion.
H	Hurst parameter
sigma	Scale parameter of l fsm
freq	Frequency of the motion. It can take two values: "H" for high frequency and "L" for the low frequency setting.
disable_X	is needed to disable computation of X. The default value is FALSE. When it is TRUE, only a levy motion is returned, which in turn reduces the computation time. The feature is particularly useful for reproducibility when combined with seeding.
levy_increments	increments of Levy motion underlying the l fsm.
seed	this parameter performs seeding of path generator

### Value

It returns a list containing the motion, the underlying Levy motion, the point number of the motions from 0 to N and the corresponding coordinate (which depends on the frequency), the parameters that were used to generate the l fsm, and the predefined frequency.

### References

Stoev S, Taqqu M (2004). “Simulation methods for linear fractional stable motion and FARIMA using the Fast Fourier Transform.” *Fractals*, **95**(1), 95-121.

**See Also**

[paths](#) simulates a number of lfsm sample paths.

**Examples**

```
# Path generation

m<-256; M<-600; N<-2^10-M
alpha<-1.8; H<-0.8; sigma<-0.3
seed=2

List<-path(N=N,m=m,M=M,alpha=alpha,H=H,
           sigma=sigma,freq='L',disable_X=FALSE,seed=3)

# Normalized paths
Norm_lfsm<-List[['lfsm']] / max(abs(List[['lfsm']]))

Norm_oLm<-List[['levy_motion']] / max(abs(List[['levy_motion']]))

# Visualization of the paths
plot(Norm_lfsm, col=2, type="l", ylab="coordinate")
lines(Norm_oLm, col=3)
leg.txt <- c("lfsm", "oLm")
legend("topright", legend = leg.txt, col =c(2,3), pch=1)

# Creating Levy motion
levyIncrems<-path(N=N, m=m, M=M, alpha, H, sigma, freq='L',
                    disable_X=TRUE, levy_increments=NULL, seed=seed)

# Creating lfsm based on the levy motion
lfsm_full<-path(m=m, M=M, alpha=alpha,
                  H=H, sigma=sigma, freq='L',
                  disable_X=FALSE,
                  levy_increments=levyIncrems$levy_increments,
                  seed=seed)

sum(levyIncrems$levy_increments ==
    lfsm_full$levy_increments)==length(lfsm_full$levy_increments)
```

paths

*Generator of a set of lfsm paths.*

**Description**

It is essentially a wrapper for [path](#) generator, which exploits the latest to create a matrix with paths in its columns.

**Usage**

```
paths(N_var, parallel, seed_list = rep(x = NULL, times = N_var), ...)
```

**Arguments**

N_var	number of l fsm paths to generate
parallel	a TRUE/FALSE flag which determines if the paths will be created in parallel or sequentially
seed_list	a numerical vector of seeds to pass to <a href="#">path</a>
...	arguments to pass to path

**See Also**

[path](#)

**Examples**

```
m<-45; M<-60; N<-2^10-M
alpha<-1.8; H<-0.8; sigma<-0.3
freq='L'
r=1; k=2; p=0.4

Y<-paths(N_var=10,parallel=TRUE,N=N,m=m,M=M,
          alpha=alpha,H=H,sigma=sigma,freq='L',
          disable_X=FALSE,levy_increments=NULL)

Hs<-apply(Y,MARGIN=2,H_hat,p=p,k=k)
hist(Hs)
```

**Description**

The function takes a list of parameters (alpha, H) and uses [expand.grid](#) to obtain all possible combinations of them. Based on each combination, the function simulates an l fsm sample path. It is meant to be used in conjunction with function [Plot\\_list\\_paths](#).

**Usage**

```
Path_array(N, m, M, l, sigma)
```

**Arguments**

N	a number of points of the l fsm.
m	discretization. A number of points between two nearby motion points
M	truncation parameter. A number of points at which the integral representing the definition of l fsm is calculated. So, after M points back we consider the rest of the integral to be 0.
l	a list of parameters to expand
sigma	Scale parameter of l fsm

**Value**

The returned value is a data frame containing paths and the corresponding values of alpha, H and frequency.

**Examples**

```
l=list(H=c(0.2,0.8),alpha=c(1,1.8), freq="H")
arr<-Path_array(N=300,m=30,M=100,l=l,sigma=0.3)
str(arr)
head(arr)
```

phi

*Phi***Description**

Defined as

$$\varphi_{\text{high}}(t; H, k)_n := V_{\text{high}}(\psi_t; k)_n \quad \text{and} \quad \varphi_{\text{low}}(t; k)_n := V_{\text{low}}(\psi_t; k)_n$$

, where  $\psi_t(x) := \cos(tx)$

**Usage**

```
phi(t, k, path, H, freq)
```

**Arguments**

t	positive real number
k	increment order
path	sample path of l fsm on which the inference is to be performed
H	Hurst parameter
freq	Frequency of the motion. It can take two values: "H" for high frequency and "L" for the low frequency setting.

## Details

Hurst parameter is required only in high frequency case. In the low frequency, there is no need to assign H a value because it will not be evaluated.

## References

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

phi_of_alpha	<i>Inverse alpha estimator</i>
--------------	--------------------------------

## Description

A function from a general estimation procedure which is defined as  $m^p - p'_k / m^{p'} - p_k$ , originally proposed in [13].

## Usage

```
phi_of_alpha(p, p_prime, alpha)
```

## Arguments

p	power
p_prime	power
alpha	self-similarity parameter of alpha stable random motion.

## References

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

Plot_dens	<i>(alpha,H,sigma)- density plot</i>
-----------	--------------------------------------

## Description

Plots the densities of the parameters (alpha,H,sigma) estimated in Monte-Carlo experiment. Works in conjunction with [MCestimLFSM](#) function.

## Usage

```
Plot_dens(par_vec = c("alpha", "H", "sigma"), MC_data, Nnorm = 1e+07)
```

**Arguments**

par_vec	vector of parameters which are to be plotted
MC_data	a list created by <a href="#">MCestimLFSM</a>
Nnorm	number of point sampled from standard normal distribution

**See Also**

[Plot\\_vb](#) to plot variance- and bias dependencies on n.

**Examples**

```
m<-45; M<-60

p<-.4; p_prime<-.2
t1<-1; t2<-2; k<-2

NmonteC<-5e2
S<-c(1e3,1e4)
alpha<-.8; H<-0.8; sigma<-0.3
theor_4_1_clt_new<-MCestimLFSM(s=S,fr='L',Nm=1e7,
                                  m=m,M=M,
                                  alpha=alpha,H=H,sigma=sigma,
                                  GenLowEstim,t1=t1,t2=t2,p=p)
l_plot<-Plot_dens(par_vec=c('sigma','alpha','H'), MC_data=theor_4_1_clt_new, Nnorm=1e7)
l_plot
```

Plot_list_paths	<i>Rendering of path lattice</i>
-----------------	----------------------------------

**Description**

Rendering of path lattice

**Usage**

```
Plot_list_paths(arr)
```

**Arguments**

arr	a data frame produced by <a href="#">Path_array</a> .
-----	---

**Examples**

```
l=list(H=c(0.2,0.8),alpha=c(1,1.8), freq="H")
arr<-Path_array(N=300,m=30,M=100,l=l,sigma=0.3)
p<-Plot_list_paths(arr)
p
```

<b>Plot_vb</b>	A function to plot variance- and bias dependencies of estimators on the lengths of sample paths. Works in conjunction with <a href="#">MCestimLFSM</a> function.
----------------	--

**Description**

A function to plot variance- and bias dependencies of estimators on the lengths of sample paths.  
Works in conjunction with [MCestimLFSM](#) function.

**Usage**

```
Plot_vb(data)
```

**Arguments**

data	a list created by <a href="#">MCestimLFSM</a>
------	---

**Value**

The function returns a ggplot2 graph.

**See Also**

[Plot\\_dens](#)

**Examples**

```
# Light weight computations

m<-25; M<-50
alpha<-1.8; H<-0.8; sigma<-0.3
S<-c(1:3)*1e2
p<-.4; p_prime<-.2; t1<-1; t2<-2
k<-2; NmonteC<-50

# Here is the continuous H-1/alpha inference procedure
theor_3_1_H_clt<-MCestimLFSM(s=S,fr='H',Nmc=NmonteC,
                                m=m,M=M,alpha=alpha,H=H,
                                sigma=sigma,ContinEstim,
                                t1=t1,t2=t2,p=p,k=k)
Plot_vb(theor_3_1_H_clt)

# More demanding example (it is better to use multicore setup)
# General low frequency inference

m<-45; M<-60
```

```

alpha<-0.8; H<-0.8; sigma<-0.3
S<-c(1:15)*1e2
p<-.4; t1<-1; t2<-2
NmonteC<-50

# Here is the continuous H-1/alpha inference procedure
theor_4_1_H_clt<-MCestimLFSM(s=S,fr='H',Nmc=NmonteC,
                                m=m,M=M,alpha=alpha,H=H,
                                sigma=sigma,GenLowEstim,
                                t1=t1,t2=t2,p=p)
Plot_vb(theor_4_1_H_clt)

```

**Retrieve\_stats***Retrieve statistics(bias, variance) of estimators based on a set of paths***Description**

Retrieve statistics(bias, variance) of estimators based on a set of paths

**Usage**

```
Retrieve_stats(paths, true_val, Est, ...)
```

**Arguments**

paths	real-valued matrix representing sample paths of the stochastic process being studied
true_val	true value of the estimated parameter
Est	estimator (i.e. H_hat)
...	parameters to pass to Est

**Examples**

```

m<-45; M<-60; N<-2^10-M
alpha<-1.8; H<-0.8; sigma<-0.3
freq='L';t1=1; t2=2
r=1; k=2; p=0.4

Y<-paths(N_var=10,parallel=TRUE,N=N,m=m,M=M,
          alpha=alpha,H=H,sigma=sigma,freq='L',
          disable_X=FALSE,levy_increments=NULL)

Retrieve_stats(paths=Y,true_val=sigma,Est=sigma_hat,t1=t1,k=2,alpha=alpha,H=H,freq="L")

```

---

R_hl	<i>R high /low</i>
------	--------------------

---

### Description

Defined as

$$R_{\text{high}}(p, k)_n := \frac{\sum_{i=2k}^n |\Delta_{i,k}^{n,2} X|^p}{\sum_{i=k}^n |\Delta_{i,k}^{n,1} X|^p},$$

$$R_{\text{low}}(p, k)_n := \frac{\sum_{i=2k}^n |\Delta_{i,k}^2 X|^p}{\sum_{i=k}^n |\Delta_{i,k}^1 X|^p}$$

### Usage

```
R_hl(p, k, path)
```

### Arguments

p	power
k	increment order
path	sample path of l fsm on which the inference is to be performed

### Details

The computation procedure for high- and low frequency cases is the same, since there is no way to control frequency given a sample path.

### References

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

### Examples

```
m<-45; M<-60; N<-2^10-M
alpha<-0.8; H<-0.8; sigma<-0.3
p<-0.3; k=3

l fsm<-path(N=N,m=m,M=M,alpha=alpha,H=H,
            sigma=sigma,freq='L',disable_X=FALSE,seed=3)$l fsm
R_hl(p=p,k=k,path=l fsm)
```

---

sf	<i>Statistic V</i>
----	--------------------

---

## Description

Statistic of the form

$$V_{\text{high}}(f; k, r)_n := \frac{1}{n} \sum_{i=rk}^n f \left( n^H \Delta_{i,k}^{n,r} X \right),$$

$$V_{\text{low}}(f; k, r)_n := \frac{1}{n} \sum_{i=rk}^n f \left( \Delta_{i,k}^r X \right)$$

## Usage

```
sf(path, f, k, r, H, freq, ...)
```

## Arguments

path	sample path for which the statistic is to be calculated.
f	function applied to high order increments.
k	order of the increments.
r	step of high order increments.
H	Hurst parameter.
freq	frequency.
...	parameters to pass to function f

## Details

Hurst parameter is required only in high frequency case. In the low frequency, there is no need to assign H a value because it will not be evaluated.

## References

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

## See Also

[phi](#) computes V statistic with  $f(\cdot) = \cos(t)$

### Examples

```

m<-45; M<-60; N<-2^10-M
alpha<-1.8; H<-0.8; sigma<-0.3
freq='L'
r=1; k=2; p=0.4
S<-(1:20)*100

path_lfsm<-function(...){

  List<-path(...)
  List$lfsm

}

Pths<-lapply(X=S,FUN=path_lfsm,
              m=m, M=M, alpha=alpha, sigma=sigma, H=H,
              freq=freq, disable_X = FALSE,
              levy_increments = NULL, seed = NULL)

f_phi<-function(t,x) cos(t*x)
f_pow<-function(x,p) (abs(x))^p

V_cos<-sapply(Pths,FUN=sf,f=f_phi,k=k,r=r,H=H,freq=freq,t=1)
ex<-exp(-(abs(sigma*Norm_alpha(h_kr,alpha=alpha,k=k,r=r,H=H,l=0)$result)^alpha))

# Illustration of the law of large numbers for phi:
plot(y=V_cos, x=S, ylim = c(0,max(V_cos)+0.1))
abline(h=ex, col='brown')

# Illustration of the law of large numbers for power functions:
Mpk<-m_pk(k=k, p=p, alpha=alpha, H=H, sigma=sigma)

sf_mod<-function(Xpath,...) {

  Path<-unlist(Xpath)
  sf(path=Path,...)
}

V_pow<-sapply(Pths,FUN=sf_mod,f=f_pow,k=k,r=r,H=H,freq=freq,p=p)
plot(y=V_pow, x=S, ylim = c(0,max(V_pow)+0.1))
abline(h=Mpk, col='brown')

```

*sigma\_hat*

*Statistical estimator for sigma*

### Description

Statistical estimator for sigma

**Usage**

```
sigma_hat(t1, k, path, alpha, H, freq)
```

**Arguments**

t1	real number such that t2 > t1 > 0
k	increment order
path	sample path of l fsm on which the inference is to be performed
alpha	self-similarity parameter of alpha stable random motion.
H	Hurst parameter
freq	Frequency of the motion. It can take two values: "H" for high frequency and "L" for the low frequency setting.

**Examples**

```
m<-45; M<-60; N<-2^14-M
alpha<-1.8; H<-0.8; sigma<-0.3
freq='H'
r=1; k=2; p=0.4; t1=1; t2=2

# Reproducing the work of ContinEstim
# in high frequency case
l fsm<-path(N=N,m=m,M=M,alpha=alpha,H=H,
             sigma=sigma,freq='L',disable_X=FALSE,seed=1)$l fsm

H_est<-H_hat(p=p,k=k,path=l fsm)
H_est
alpha_est<-alpha_hat(t1=t1,t2=t2,k=k,path=l fsm,H=H_est,freq=freq)
alpha_est

sigma_est<-tryCatch(
  sigma_hat(t1=t1,k=k,path=l fsm,
            alpha=alpha_est,H=H_est,freq=freq),
  error=function(c) 'Impossible to compute sigma_est')
sigma_est
```

$$\theta(g, h)_p = a_p^{-2} - 2 \int_{\mathbb{R}^2} |xy|^{-1-p} U_{g,h}(x, y) dx dy$$

**Description**

Function of the form

$$\theta(g, h)_p = a_p^{-2} \int_{\mathbb{R}^2} |xy|^{-1-p} U_{g,h}(x, y) dx dy$$

**Usage**

```
theta(p, alpha, sigma, g, h)
```

**Arguments**

<i>p</i>	power, real number from (-1,1)
<i>alpha</i>	self-similarity parameter of alpha stable random motion.
<i>sigma</i>	Scale parameter of Ifsm
<i>g</i>	functions $g, h : \mathbb{R} \rightarrow \mathbb{R}$ with finite alpha-norm (see <a href="#">Norm_alpha</a> ).
<i>h</i>	functions $g, h : \mathbb{R} \rightarrow \mathbb{R}$ with finite alpha-norm (see <a href="#">Norm_alpha</a> ).

**References**

Mazur S, Otryakhin D, Podolskij M (2018). “Estimation of the linear fractional stable motion.” Aarhus University.

*U\_g**alpha norm of u\*g***Description**

alpha norm of  $u^*g$

**Usage**

```
U_g(g, u, ...)
```

**Arguments**

<i>g</i>	functions $g, h : \mathbb{R} \rightarrow \mathbb{R}$ with finite alpha-norm (see <a href="#">Norm_alpha</a> ).
<i>u</i>	real numbers
...	additional parameters to pass to Norm_alpha

**Examples**

```
g<-function(x) exp(-x^2)
g<-function(x) exp(-abs(x))
U_g(g=g,u=4,alpha=1.7)
```

---

U_gh	<i>alpha-norm of u*g + v*h.</i>
------	---------------------------------

---

**Description**

alpha-norm of  $u^*g + v^*h$ .

**Usage**

`U_gh(g, h, u, v, ...)`

**Arguments**

g	functions $g, h : \mathbb{R} \rightarrow \mathbb{R}$ with finite alpha-norm (see <a href="#">Norm_alpha</a> ).
h	functions $g, h : \mathbb{R} \rightarrow \mathbb{R}$ with finite alpha-norm (see <a href="#">Norm_alpha</a> ).
u	real numbers
v	real numbers
...	additional parameters to pass to Norm_alpha

**Examples**

```
g<-function(x) exp(-x^2)
h<-function(x) exp(-abs(x))
U_gh(g=g, h=h, u=4, v=3, alpha=1.7)
```

---

U_ghuv	<i>A dependence structure of 2 random variables.</i>
--------	--

---

**Description**

It is used when random variables do not have finite second moments, and thus, the covariance matrix is not defined. For  $X = \int_{\mathbb{R}} g_s dL_s$  and  $Y = \int_{\mathbb{R}} h_s dL_s$  with  $\|g\|_{\alpha}, \|h\|_{\alpha} < \infty$ . Then the measure of dependence is given by  $U_{g,h} : \mathbb{R}^2 \rightarrow \mathbb{R}$  via

$$U_{g,h}(u, v) = \exp(-\sigma^{\alpha} \|ug + vh\|_{\alpha}^{\alpha}) - \exp(-\sigma^{\alpha} (\|ug\|_{\alpha}^{\alpha} + \|vh\|_{\alpha}^{\alpha}))$$

**Usage**

`U_ghuv(alpha, sigma, g, h, u, v, ...)`

**Arguments**

alpha	self-similarity parameter of alpha stable random motion.
sigma	Scale parameter of Ifsm
g, h	functions $g, h : \mathbb{R} \rightarrow \mathbb{R}$ with finite alpha-norm (see <a href="#">Norm_alpha</a> ).
v, u	real numbers
...	additional parameters to pass to U_gh and U_g

**Examples**

```
g<-function(x) exp(-x^2)
h<-function(x) exp(-abs(x))
U_ghuv(alpha=1.5, sigma=1, g=g, h=h, u=10, v=15,
rel.tol = .Machine$double.eps^0.25, abs.tol=1e-11)
```

# Index

a\_p, 4  
a\_tilda, 4  
alpha\_hat, 2  
  
CLT, 5  
ContinEstim, 6  
  
expand.grid, 18  
  
GenHighEstim, 7  
GenLowEstim, 8  
  
H\_hat, 9, 13, 14  
h\_kr, 9  
  
increment, 10, 14  
increments (increment), 10  
  
m\_pk, 14  
MCestimLFSM, 5, 11, 20–22  
MinContrastEstim, 13  
  
Norm\_alpha, 15, 28, 29  
  
path, 16, 17, 18  
Path\_array, 18, 21  
paths, 17, 17  
phi, 3, 14, 19, 25  
phi\_of\_alpha, 20  
Plot\_dens, 20, 22  
Plot\_list\_paths, 18, 21  
Plot\_vb, 21, 22  
  
R\_hl, 24  
Retrieve\_stats, 23  
  
sf, 25  
sigma\_hat, 26  
  
theta, 27  
  
U\_g, 28  
U\_gh, 29  
U\_ghuv, 29  
uniroot, 7